

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY
WASHINGTON, D.C. 20242

Technical Letter
NASA - 61
November 1966

Dr. Peter C. Badgley
Chief, National Resources Program
Office of Space Science and Applications
Code SAR, NASA Headquarters
Washington, D.C. 20546

Dear Peter:

Transmitted herewith are 3 copies of

TECHNICAL LETTER NASA-61
HYDROLOGIC INTERPRETATION OF
NIMBUS VIDICON IMAGE -- GREAT SALT LAKE, UTAH*

by

D. C. Hahl and A. H. Handy**

Sincerely yours,

W. A. Fischer

William A. Fischer
Research Coordinator
Earth Orbiter Program

**U.S. Geological Survey, Water Resources Division, Salt Lake City, Utah.
*Work performed under NASA contract R-09-020-015

N70-38843

FACILITY FORM 602

(ACCESSION NUMBER)	(THRU)
6	
(PAGES)	(CODE)
AR-79887	13
(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

U. S. Government Agencies Only

R-09-020-015

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

TECHNICAL LETTER NASA-61
HYDROLOGIC INTERPRETATION OF
NIMBUS VIDICON IMAGE -- GREAT SALT LAKE, UTAH*

by

D. C. Hahl and A. H. Handy

November 1966

These data are preliminary and should
not be quoted without permission

Prepared by the Geological Survey for
National Aeronautics and Space
Administration (NASA)

*Work performed under NASA Contract No. R.-09-020-015

U. S. Government Agencies Only

HYDROLOGIC INTERPRETATION OF NIMBUS VIDICON IMAGE GREAT SALT LAKE, UTAH

by

D. C. Hahl and A. H. Handy

A Nimbus Advanced Vidicon Camera System (AVCS) image of Great Salt Lake and adjacent areas recorded in September 1964 was examined to determine if features of hydrologic significance were visible and identifiable. An annotated copy of the image is shown in figure 1.

Evidently several factors are required to produce an image. Features such as Willard Bay Reservoir and Utah Lake do not show on the image, yet Great Salt Lake, which is of comparable depth shows clearly. Adjoining Willard Bay Reservoir is Bear River Bay Migratory Bird Refuge, which is discernible. The dikes, land areas, and water areas in the refuge match those on recent aerial photographs. Depth does not seem to be a factor in producing an image because the brackish water of the refuge that is less than two feet deep shows black; whereas, subaqueous features ("X's" offshore of Great Salt Lake, figure 1) of comparable depth in Great Salt Lake show through the brine.

Vegetation and geology may be responsible for the delineation or lack of delineation of some features. An example is a comparison between White Valley and Floating Island. White Valley is surrounded by mountains of about equal height (4,500 feet above the valley floor), yet only the eastern mountains are visible. Floating Island is a small feature (800 feet above the desert floor) and it shows clearly in the image.

The only man-made features discernible are the Bear River Bay Migratory Bird Refuge and the open pit copper mine of Kennecott Copper Corporation at Bingham.

Hydrologic interpretations of the image must be based on the indirect evidence of geographic and geologic features seen in the image. The most recognizable feature is Great Salt Lake. Possibly additional Nimbus pictures may aid in determining the change in shape and size of the lake with change in stage even though there is some penetration of the water. Correlation of the edge of the lake as shown in the image with depth is not possible because of the coarse resolution of the image. Seasonal changes in the size and shape of desert areas may be shown by additional pictures if the shading is due to vegetation.

The AVCS image is of interest to hydrologists. Though the cause of changes in shading on the image will be recorded only by ground monitoring, remote-sensing techniques present a means of recording occurrences heretofore not visualized. In an area such as Utah where diverse geography, geology, and hydrology exist, a key to photographic tones can possibly be established that will be applicable in interpreting data from unstudied areas. The synoptic view would be helpful in planning better coordinated ground work.

NIMBUS SATELLITE AVCS IMAGERY GREAT SALT LAKE, UTAH

